



US006113810A

United States Patent [19][11] **Patent Number:** **6,113,810****Hou et al.**[45] **Date of Patent:** **Sep. 5, 2000**

[54] **METHODS OF PREPARING
ELECTROPHORETIC DISPERSIONS
CONTAINING TWO TYPES OF PARTICLES
WITH DIFFERENT COLORS AND OPPOSITE
CHARGES**

5,229,242 7/1993 Mahabadi et al. 430/106.6
5,230,978 7/1993 Kawasaki et al. 430/106

OTHER PUBLICATIONS

Colloids and Surfaces, 6 (1983) p. 83, Luckham et al., no month available.

Colloids and Surfaces, 6 (1983) p. 101, Luckham et al., no month available.

Journal of Chemical Society, Farada 1, 76 (1980), p., 655 no month available.

[75] Inventors: **Wei-Hsin Hou**, Bethlehem, Pa.;
Frederic E. Schubert, Shoreham, N.Y.

[73] Assignee: **Copytele, Inc.**, Melville, N.Y.

[21] Appl. No.: **08/383,667**

[22] Filed: **Feb. 3, 1995**

Primary Examiner—Christine Skane

Attorney, Agent, or Firm—Arthur L. Plevy; Buchanan
Ingersoll PC

Related U.S. Application Data

[63] Continuation of application No. 08/065,871, May 21, 1993, abandoned.

[51] **Int. Cl.**⁷ **H01B 3/00**

[52] **U.S. Cl.** **252/572; 313/483; 359/296**

[58] **Field of Search** 252/572, 73, 77,
252/79; 313/483; 204/299 R; 359/296

[56] **References Cited**

U.S. PATENT DOCUMENTS

3,484,162 12/1969 Clark 255/3
4,093,534 6/1978 Carter et al. 204/299 R
4,285,801 8/1981 Chiang 204/299 R
4,298,448 11/1981 Muller et al. 204/299 R
4,655,897 4/1987 DiSanto et al. 204/299 R
4,992,192 2/1991 Ahmed 252/73

[57] **ABSTRACT**

A dielectric dispersion for use in an electrophoretic display includes a dielectric fluid, a first plurality of particles of a first color having a surface charge of a selected polarity dispersed within the dielectric fluid and a second plurality of particles of a second color having a surface charge of opposite polarity to that of the first plurality and a steric repulsion thereto preventing coagulation of the first and second pluralities. In one embodiment, the first and second plurality of particles are each formed by separate two stage dispersion polymerization reactions. Each set of particles is formed with unique secondary and functional monomers. Corresponding charge control agents are added to the dispersion to establish opposite polarities on the respective particles.

18 Claims, 1 Drawing Sheet